

ORIGINAL PAPER

Dental Morphological Anomalies in the Adi Tribe of Pasighat in Arunachal Pradesh

Das Lima¹, Bhuyan AC², Kataki Rubi³, Kalita Chandana⁴

Received on October 26, 2017; *editorial approval* on November 23, 2017

ABSTRACT

Objective: To identify three dental morphological anomalies of permanent teeth namely Peg Laterals, Dens Evaginatus and Rudimentary third molars in the Adi tribe of Pasighat area of Arunachal Pradesh. **Methods:** Oral examination for morphological anomalies was done in 156 individuals using a dental mirror and probe for the said anomalies and a questionnaire with details relevant to the study was used during examination. **Results:** A total of 156 samples were screened and 35% presented with at least one anomaly out of which 31% had one anomaly, 4% had two anomalies and 0% had more than two anomalies. Total number of positive cases (both single and dual) for each of the anomalies was- Peg Laterals 25%, Dens Evaginatus 20% and Rudimentary third molar 4%. One other anomaly, i.e. Cusp of Carabelli not previously considered in this study was found in 22% of the cases. **Conclusion:** The data obtained from the present study confirms the prevalence of dental anomalies namely, Peg Laterals, Dens Evaginatus, Rudimentary third molars and Cusp of Carabelli in the tribal population of Pasighat similar to other Mongoloid populations of the world.

Keywords: Cusp, Peg Laterals, Dens Evaginatus, Rudimentary third molars, Cusp of Carabelli

INTRODUCTION

Dental morphological anomalies in permanent teeth are an important area of study within the subject of Dental Anatomy and Physiology. There are several reasons for studying dental anatomical variants in any population, the most significant being **enhanced diagnosis and treatment planning**. Early diagnosis of dental anomalies allow for more comprehensive treatment planning, better prognosis and in certain instances, less extensive interception.¹ Another important reason for studying anatomical variation is for **forensic value**. Anatomical variation related to tooth morphology may show certain traits specific to different racial groups which may

be of immense forensic value. Ancestry can be accessed by studying the facial skeleton and comparing the features with the main characteristics of three racial groups: Mongoloid, Negroid, and Caucasoid.² Also, the large variation in morphological features and their form may not be easily altered; thus a trait of the human dentition can be a valuable diagnostic tool for **anthropological studies** in classifying and characterizing different ethnic groups.³ Accumulation of data on the morphological traits of teeth in different populations has let anthropologists to research further on the evolutionary significance of this data and consider its mode of inheritance. Again the incidence and degree of expression of anomalies in different population groups can provide important information for phylogenetic and genetic studies and help the understanding of variations within and between the different world populations.⁴

However, the occurrence of dental morphologic anomalies has not been adequately researched in the remote tribes and sub-tribes of north-east India. The present study will not only contribute towards a better understanding of the said topic but will also provide a base line data for allied branches to review.

An initial hypothesis related to the present research is as follows-Three dental anomalies of permanent teeth namely Peg Lateral, Dens Evaginatus and Microdontia of third molars are endemic in the Adi population of Pasighat, in Arunachal Pradesh.

Address for correspondence:

¹Reader (**Corresponding Author**)

Email: limakaling@gmail.com

Mobile: +919954714031

²Professor and Head, ³Professor, ⁴Reader

Dept. of Conservative Dentistry & Endodontics
Regional Dental College, Guwahati, Assam

Objectives: The aim of the study is to identify three dental morphological anomalies namely Peg laterals, Dens Evaginatus and Microdontia of maxillary third molars/Rudimentary third molars (erupted) in the tribal population in Pasighat area of Arunachal Pradesh, India.

The overall objective of the proposed study was to establish through data the occurrence of dental morphological anomalies in permanent teeth in the said population.

MATERIALS AND METHODS

Oral examination was done using dental mirror and probe for the said anomalies and a questionnaire with details relevant to the study was used during examination. The age group selected was above 21 years. A stratified cluster sampling method for data collection was used after dividing the said Pasighat area into three main geographical groups and further stratifying them into sub groups. The three main groups are: Pasighat urban subdivision (gr 1), Balek group of villages (gr 2) and Mirmir group of villages.

A total number of 156 samples were screened in this present study distributed in the three main groups in the ratio 12:2:5 according to population ratio. Geographical area concerned with the study includes the entire Pasighat township within 10 km radius.

Inclusion & exclusion criteria were considered as follows:

Inclusion criteria: Persons belonging to the Adi tribe and age should be above 21 years.

Exclusion criteria: If one or both parents are not from the Adi tribe, restored teeth and unerupted.

third molars. Study variables such as sub tribes of the Adi tribe (e.g: pasi, padam, panggi, minyong, milang), gender, type of anomaly and geographical distribution were

considered.

Definitions:

Peg Laterals are maxillary lateral incisors in which the mesial and distal sides converge or taper together incisally forming a peg shaped or cone shaped crown instead of exhibiting parallel or diverging proximal surfaces. Dens Evaginatus is a developmental condition that appears clinically as an accessory cusp or a globule of enamel usually on the occlusal surface between the buccal and lingual cusps of premolars unilaterally or bilaterally. Again, microdontia of third maxillary molars/rudimentary third molar are third molar teeth that are smaller in size than normal, i.e., outside the usual limits of variations.⁵

Ethical clearance was taken from the institutional ethics committee. Data analysis was done using IBM SPSS version 20 and Microsoft Excel Spreadsheet software.

RESULTS

Oral examination showed the presence of three previously mentioned anomalies Peg laterals, Dens Evaginatus and Microdontia of third molars and also included positives of one other anomaly, i.e. Cusp of Carabelli in a number of the cases. Of the total sample of 156 the number of cases with anomalies was 55, i.e. 35% of the total cases screened. Of the total 156 samples number of Cusp of Carabelli found were 34, i.e., 22%, Peg laterals (PL) 14, i.e. 9%, Dens Evaginatus (DE) 11 i.e., 7%, Microdontia of third molars (RT) were 2 i.e., 1%. Of the positive 55 cases the number and percentage of individual anomalies (both single and dual) are Cusp of Carabelli 34, i.e., 62%, Peg laterals (PL) 14, i.e., 25%, Dens Evaginatus (DE) 11, i.e., 20%, Microdontia of third molars (RT) 2, i.e., 4%. Again, the prevalence of anomalies was 59% in the females and 41% in the males (**Figure 2**). The distributions of anomalies in the various sub-tribes were Pasi-18%, Padam - 16%, Milang - 2%, Pangi - 4% and Minyong - 60% (**Figure 3**). Geographically, 45% of the anomalies were found in the urban area, 24% in the Balek area and 31% in the Mirmir area (**Figure 4**).

Table 1 Number and percentage of occurrence of anomalies out of the total samples and among the positives

Descriptive Statistics	Observed values	% of total sample (156)	% of positive cases (55)
Total no. Of screenings (sample size)	156		
No. Of positive cases	55	35%	
No. Of cases with one anomaly	49	31%	89%
No. Of cases with two anomalies	6	4%	11%
No. Of cases with more than two anomalies	0	0%	0%
No. Of pl positive cases	14	9%	25%
No. Of de positive cases	11	7%	20%
No. Of rt positive cases	2	1%	4%
No. Of cc positive cases	34	22%	62%

**TOTAL CASES OF PL, DE, RT & CC (SINGLE + DUAL)
AS A % OF THE TOTAL SAMPLE**




Figure 1 Distribution of anomalies as a percentage of the total sample

GENDER WISE DISTRIBUTION OF THE SAMPLE




Figure 2 Gender-wise distribution of anomalies

**SUB TRIBE WISE DISTRIBUTION OF THE POSITIVE
CASES**




Figure 3 Distribution of anomalies according to the sub-tribes

**GEOGRAPHICAL AREA WISE DISTRIBUTION OF THE
SAMPLE**




Figure 4 Geographical distributions of anomalies

DISCUSSION

The occurrence of dental morphological anomalies has been studied by various authors in different world populations at various times. However such studies identifying dental morphological traits in the various tribes and sub-tribes of the north-eastern states of India have not received its due importance.^{3, 6, 7}

In the present study, of the total number of 156 cases, 35% presented with at least one anomaly. 31% of the total cases had one anomaly, 4% had two anomalies and none had more than two anomalies. It was observed that 9% of the total cases were positive for Peg Lateral, 7% for Dens Evaginatus, 1% for rudimentary third molar and 22% for Cusp of Carabelli. Again to statistically analyze the results A 'Z' test of proportion was done to test the difference between the hypothesized proportion of anomalies ($P=50\%$) and the observed proportion ($p=35\%$) at 5% significance level. The Z test indicated that the p value < 0.05 .

Chi square tests were conducted at 5 % significance level to test the difference in the distribution of anomalies according to gender, area and the various sub tribes. The tests indicated that there is no significant difference in the distribution of anomalies with

respect to gender (p value = 0.85). However, highly significant differences were indicated in the distribution of the anomalies with respect to area (p value < 0.001) and to sub tribe (p value < 0.001).

A study similar to our present study was done by Nayak P in Rajasthan and he found that prevalence of peg lateral was 0.4% which is very less compared to the 9% prevalence in our study.⁸ Hua F also did a meta analysis on the prevalence of peg-shaped maxillary permanent lateral incisors and their associations with race, population type, sex, and sidedness.⁹ The overall prevalence of peg-shaped maxillary permanent lateral incisors was 1.8%, the occurrence rates being higher in Mongoloid (3.1%) than in black (1.5%) and white (1.3%) patients. Comparatively the findings in our study showed a higher percentage of prevalence of Peg laterals. In his study women were more likely than men to have peg shaped maxillary permanent lateral incisors. The prevalence rates of unilateral and bilateral peg-shaped maxillary permanent lateral incisors were approximately the same. However, among the unilateral lateral incisors, the left side (0.4%) was twice as common as the rightside (0.2%). Another similar study on the dental features such as size, shape, cusp number and groove

pattern, etc. was done on the Tibetan immigrants of India by Sharma J.¹⁰ He concluded that overall reduction in size, hypodontia of the third molar and absence of Carabelli's Cusp were a distinct evolutionary trend in Tibetan dentition. In our study however the prevalence of the Cusp of Carabelli was much higher, i.e. 22%. In another study by Stecker SS the prevalence of dental anomalies in a Southeast Asian Population in the Minneapolis area reported a higher 7.5% prevalence of peg laterals and 1.1% of Dens Evaginatus in Asians.¹¹

In our study since no previous data relating to the prevalence of dental morphological anomalies in the Adi tribe was found, population proportion (P) with anomalies was hypothetically taken as 50%. The observed proportion following data collection and analysis was 35%. The Z test used to compare the two proportions rejected the null hypothesis. However the higher prevalence rate confirmed the presence of specific types of dental anomalies and has given us their prevalence rates for future studies.

CONCLUSION

The data obtained from the present study confirms the prevalence of dental anomalies, namely Peg Lateral, Dens Evaginatus, Microdontia of third molars and Cusp of Carabelli in the Pasighat area of Arunachal Pradesh

Conflict of interest: No conflict of interest associated with this work.

Ethical clearance: Taken.

Source of funding: None declared.

Author declaration: We declare that this work was done by the authors named in this article and all liabilities pertaining to claims relating to the content of this article will be borne by the authors.

REFERENCES

1. Yonezu T, Hayashi Y, Sasaki J, Machida Y. Prevalence of congenital dental anomalies of the deciduous dentition in Japanese children. *Bull Tokyo DentColl* 1997;38:27-32.
2. Leung KK. Forensic odontology. *Dent Bull* 2008;13:11.
3. King N M, Tsai Jennie S J, Wong H M. Morphological and numerical characteristics of the southern chinese dentitions. Part II: Traits in the permanent dentition. *Open Anthropol J* 2010;3:71-84.
4. Patil S, Doni B, Kaswan S, Rahman F. Prevalence of dental anomalies in Indian population. 2013 Oct. [cited 2014 Jul];5(4): [e183–e186]. Available from: URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3892239/>
5. Shafer WG, Hine MK, Levy BM. *Oral Pathology*. 5thed. Singapore: Harcourt Brace Asia; 1993.p. 52-59.
6. Jerome CE, Hanlon RJ. Dental anatomical anomalies in asians and pacific islanders. *CDA J* 2007;35(9):631-6.
7. Lavelle CLB. A metrical comparison of maxillary first premolar form. *Am J Phys Anthropol* 1984;63:397-403.
8. Nayak P, Nayak S. Prevalence and distribution of dental anomalies in 500 Indian school children. *Bang J of med Sc* 2011;10(1).
9. Hua F, He H, Ngan P. Prevalence of peg-shaped maxillary permanent lateral incisors: a meta-analysis. *Am J of Orthod and Dentofacial Orthop* 2013;144(1):97-109.
10. Sharma J. Dental morphology and odontometry of the tibetan immigrants. *Am J PhysAnthropol* 1983;61(4):495-505.
11. Stecker SS, Beirahi S, Hodqes JS, Peterson VS. Prevalence of dental anomalies in a southeast asian population in the minneapolis/saint paul metropolitan area. *J of Minnesota dental assoc* 2013;92(4).

ORIGINAL PAPER

Pattern of Homicide in Gurgaon Region

Chauhan Harsh¹, Yadav Ruchika², Mathur Deepak³

Received on August 02, 2017; editorial approval on September 01, 2017

ABSTRACT

Introduction: Homicide means killing of human being. It is very heinous crime done by a person against other human being. **Aim:** Aim of this study was to find out the most prominent method of homicide in Gurgaon region. **Method:** This study was a retrospective study done in the Department of Forensic Medicine & Toxicology, Faculty of Medicine & Health Sciences, SGT University, Gurgaon in association with Mortuary of Civil Hospital Gurgaon. **Results:** A total of 1196 autopsies were conducted at Mortuary, Civil hospital Gurgaon during the period from January 2016 to December 2016, out of which, 60(5%) cases were of alleged homicidal death. Majority of victims were in the age group of 31-40 years with Male:Female ratio; 4.1 were observed. 78.3% victims were male. 51.67% of total victims were resident of place outside Gurgaon with majority of male victims. Contusion was found in maximum cases followed by fracture 78.3% death were due to mechanical injuries followed by asphyxial death which is in 21.7%. Head injury was observed as a major cause of death in 36.7% cases. **Conclusion:** The most affecting age group was 31-40 years with male predominance. 78.33% cases were died due to mechanical injuries, out of which maximum due to blunt weapon.

Keywords: Sharp weapon, blunt weapon, firearm

INTRODUCTION

Life and death are two undeniable facts in this world. Like birth, death is also inevitable. There is only one way for birth, there are many ways to die. Homicide is legally defined as destruction of human life by the act, agencies, procurement or culpable omission of some other person or persons.¹ Homicide is prevalent all over the world.¹ Globally around 5,20,000 people die every year due to interpersonal violence, which equates nearly 1400 deaths every day.

The various methods of homicidal deaths are injury by blunt weapon, sharp weapon, firearm, strangulation, homicidal hanging (Lynching), smothering, poisoning etc. The pattern and incidences of homicides are increasing because of

population explosion, changes in life style, modern needs of men and easy availability of various types of weapons. In view of the magnitude and frequency of such deaths and its impact on the society, the present study is under taken so as to find out the most vulnerable age group, sex incidence, pattern of homicide, residence of victim of crime, region of the body involved and cause of death.

In India, after many years of independence, the rate of homicides is increasing day by day. As per National Crime Record Bureau, violent crimes reported in India were 10.9% of the total Indian Penal Code crimes. The total number of murders recorded all over the India in 2010 was 33,335. There are consistent differences in rates of homicide victimization between males and females and young and old. In terms of age difference, homicide victimization rates are generally higher for young adults, especially young adult males.

MATERIALS AND METHODS

A retrospective study regarding the various patterns of homicide in Gurgaon region was done in the Department of Forensic Medicine & Toxicology, Faculty of Medicine & Health Sciences, SGT University, Gurgaon in association with Mortuary of Civil Hospital Gurgaon during the period from January 2016 to December 2016. All the cases were included in this study who were brought to the mortuary of Civil Hospital Gurgaon for the medico-legal post-mortem examination with alleged history of homicide. Cases of unnatural deaths where the cause of death could not be

Address for correspondence:

¹Post-graduate student (**Corresponding Author**)

SGT Medical College, Hospital & Research Institute, Budhera, Gurgaon

Email: forensicmedicine2018@gmail.com

Mobile: +918375800981

²Assistant Professor of Forensic Science

SGT Medical College, Hospital & Research Institute, Budhera, Gurgaon

³Forensic Expert, Civil Hospital, Gurgaon.